

Care Tracker

This Care Tracker is meant to provide you and your healthcare team with a more detailed understanding of how DAYBUE® or DAYBUE® STIX is affecting the signs and symptoms of Rett syndrome in your loved one. Use the first sheet to note the range of symptoms, behaviors, and moods you've seen in your loved one before treatment, then regularly fill out the rest to track any changes over time.

How to use:

- ✓ Capture the day-to-day changes you notice in your loved one with DAYBUE or DAYBUE STIX
- ✓ Ask people in your loved one's care circle—like family, friends, teachers, and therapists—about any changes they've observed
- ✓ Note any changes in dosage or diet
- ✓ Note any side effects that you have noticed in your loved one. Tell your healthcare provider if your loved one has any side effects that are bothersome or do not go away
- ✓ Start a new sheet weekly, monthly, or at your own pace
- ✓ Share with your loved one's doctor

Save, scan, or take a picture of each completed sheet and ask your doctor to consider saving a copy with your loved one's records.

Visit [DAYBUE.com](https://www.daybue.com) for more information about DAYBUE and DAYBUE STIX and remember that your Acadia Connect® Family Support Educator is available to help you and your loved one in their Rett syndrome journey.

Indication

What is DAYBUE or DAYBUE STIX?

DAYBUE or DAYBUE STIX are prescription medicines used to treat Rett syndrome in adults and children 2 years of age and older. It is not known if DAYBUE or DAYBUE STIX is safe and effective in children under 2 years of age.

Important Safety Information

What are the possible side effects of DAYBUE or DAYBUE STIX?

- **Diarrhea:** Diarrhea is a common side effect of DAYBUE or DAYBUE STIX that can sometimes be severe. Diarrhea may cause you to lose too much water from your body (dehydration). Before starting treatment with DAYBUE or DAYBUE STIX, stop taking laxatives. Tell your healthcare provider if you have diarrhea while taking DAYBUE or DAYBUE STIX. Your healthcare provider may ask you to increase the amount you drink or take antidiarrheal medicine as needed.

See additional Important Safety Information on [next page](#). Please read the full [Prescribing Information](#), including [Patient Information](#).



Indication and Important Safety Information

What is DAYBUE or DAYBUE STIX?

DAYBUE or DAYBUE STIX are prescription medicines used to treat Rett syndrome in adults and children 2 years of age and older. It is not known if DAYBUE or DAYBUE STIX is safe and effective in children under 2 years of age.

What are the possible side effects of DAYBUE or DAYBUE STIX?

- **Diarrhea:** Diarrhea is a common side effect of DAYBUE or DAYBUE STIX that can sometimes be severe. Diarrhea may cause you to lose too much water from your body (dehydration). Before starting treatment with DAYBUE or DAYBUE STIX, stop taking laxatives. Tell your healthcare provider if you have diarrhea while taking DAYBUE or DAYBUE STIX. Your healthcare provider may ask you to increase the amount you drink or take antidiarrheal medicine as needed.
- **Vomiting:** Vomiting is a common side effect of DAYBUE or DAYBUE STIX. Sometimes vomit can get into your lungs (aspiration), which could cause an infection (aspiration pneumonia). Tell your healthcare provider if you have severe vomiting or if vomiting happens often.
- **Weight loss:** DAYBUE or DAYBUE STIX can cause weight loss. Tell your healthcare provider if you notice you are losing weight at any time during treatment with DAYBUE or DAYBUE STIX.
- **The most common side effects of DAYBUE or DAYBUE STIX were diarrhea and vomiting.** Other side effects include fever, seizure, anxiety, decreased appetite, tiredness, and the common cold. These are not all the possible side effects of DAYBUE or DAYBUE STIX. Tell your healthcare provider if you have any side effects that bother you or do not go away.

What should I tell my healthcare provider before taking DAYBUE or DAYBUE STIX?

Before taking DAYBUE or DAYBUE STIX, tell your healthcare provider about all of your medical conditions, including if you:

- have kidney problems.
- are pregnant or plan to become pregnant. It is not known if DAYBUE or DAYBUE STIX will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if DAYBUE or DAYBUE STIX passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby while taking DAYBUE or DAYBUE STIX.

Tell your healthcare provider about all of the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking DAYBUE or DAYBUE STIX with certain medicines may affect how the other medicines work and can cause serious side effects.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call **1-800-FDA-1088**. You can also call Acadia Pharmaceuticals Inc. at **1-844-4ACADIA (1-844-422-2342)**.

DAYBUE is available as an oral solution (200 mg/mL).

DAYBUE STIX for oral solution powder is available in 5,000 mg, 6,000 mg, and 8,000 mg packets.

The risk information provided here is not comprehensive. To learn more, talk with your healthcare provider and read the full [Prescribing Information](#), including [Patient Information](#), also available at DAYBUE.com.



Care Tracker

Name: _____

Date: _____

Use this tracker to record any changes you notice while your loved one is taking DAYBUE or DAYBUE STIX. Even small details may give your doctor valuable insight into how your loved one is responding to DAYBUE or DAYBUE STIX. **Save, scan, or take a picture of each completed sheet and ask your doctor to consider saving a copy with your loved one's records.**

Use the spaces below and on the back to track changes. The descriptions provide examples of what to look for.

 Breathing Hyperventilating, breath holding, or "blue spells"	<i>(eg, Fewer "blue spells")</i>
 Hand movements or stereotypies Restricted use of hands or involuntary, repetitive, or patterned movements	<i>(eg, Hands are more fluid and open)</i>
 Repetitive behaviors May include repetitive tongue or hand movements, rocking back and forth	<i>(eg, Teeth grinding much less)</i>
 Nighttime behaviors Crying, laughing, or screaming at night for no reason at all	<i>(eg, Fewer night terrors)</i>
 Vocalizations Uses nonverbal or verbal communication to express needs or wants	<i>(eg, Greater use of sounds as a response)</i>
 Facial expressions Grimacing expression with face or mouth or lack of facial expression	<i>(eg, Doesn't stick tongue out as much)</i>
 Eye gaze Uses eye gaze to convey feelings, needs, or wishes	<i>(eg, Can hold gaze to participate and engage)</i>
 Mood Crying, screaming, or irritability for no apparent reason	<i>(eg, Mood is better at home, in school, and in groups)</i>

See Important Safety Information on page 2. Please read the full [Prescribing Information](#), including [Patient Information](#).



Observations from other physicians, therapists, teachers, friends, or family members:

Describe side effects:

Tell your healthcare provider if your loved one has any side effects that are new, bothersome, or do not go away.

Dietary changes:

Additional notes, observations, or questions you may have:

See Important Safety Information on page 2. Please read the full [Prescribing Information](#), including [Patient Information](#).

