


Talking with your doctor
about DAYBUE® (trofinetide)

Every small
improvement
with DAYBUE may
open up more
possibilities

DAYBUE is the first and only
FDA-approved treatment
for Rett syndrome

MEET DAYBUE® STIX

Now families have a choice



*Maddy, age 21, on DAYBUE
since 2020, with her sister*

Indication

What is DAYBUE or DAYBUE STIX?

DAYBUE or DAYBUE STIX are prescription medicines used to treat Rett syndrome in adults and children 2 years of age and older. It is not known if DAYBUE or DAYBUE STIX is safe and effective in children under 2 years of age.

Important Safety Information

What are the possible side effects of DAYBUE or DAYBUE STIX?

- **Diarrhea:** Diarrhea is a common side effect of DAYBUE or DAYBUE STIX that can sometimes be severe. Diarrhea may cause you to lose too much water from your body (dehydration). Before starting treatment with DAYBUE or DAYBUE STIX, stop taking laxatives. Tell your healthcare provider if you have diarrhea while taking DAYBUE or DAYBUE STIX. Your healthcare provider may ask you to increase the amount you drink or take antidiarrheal medicine as needed.

See additional Important Safety Information on page 6.

Please read the full [Prescribing Information](#), including [Patient Information](#).





Talking to your doctor about DAYBUE

Rett syndrome is a complex neurodevelopmental disorder that is typically caused by a mutation, or change, in the *MECP2* gene. DAYBUE is the first and only FDA-approved treatment for adults and children ages 2 and older with Rett syndrome.

With the FDA approval of a treatment for Rett syndrome, you may be wondering if DAYBUE or DAYBUE STIX is right for your loved one.

In this guide, you'll find helpful information about DAYBUE and some questions to help you prepare for a conversation with your loved one's doctor.



Talk to your doctor to find out more about DAYBUE and whether it could be right for your loved one.

Visit [DAYBUE.com](https://www.daybue.com) to learn more.

Important Safety Information (continued)

What are the possible side effects of DAYBUE or DAYBUE STIX? (continued)

- **Vomiting:** Vomiting is a common side effect of DAYBUE or DAYBUE STIX. Sometimes vomit can get into your lungs (aspiration), which could cause an infection (aspiration pneumonia). Tell your healthcare provider if you have severe vomiting or if vomiting happens often.
- **Weight loss:** DAYBUE or DAYBUE STIX can cause weight loss. Tell your healthcare provider if you notice you are losing weight at any time during treatment with DAYBUE or DAYBUE STIX.
- **The most common side effects of DAYBUE or DAYBUE STIX include diarrhea and vomiting.** Other side effects include fever, seizure, anxiety, decreased appetite, tiredness, and the common cold. These are not all the possible side effects of DAYBUE or DAYBUE STIX. Tell your healthcare provider if you have any side effects that bother you or do not go away.

See additional Important Safety Information on page 6.



What to look for

- What symptoms could DAYBUE help address?
- When could I start to see improvements?
- Are improvements the same for everyone?
- Will improvements stop if my loved one stops taking DAYBUE?
- How long will my loved one need to be on DAYBUE?



Deciding if DAYBUE is right for your loved one

- Who is DAYBUE for?
- Does DAYBUE take the place of other medications my loved one is on?
- Will DAYBUE interact with other medications my loved one takes?

Important Safety Information (continued)

What should I tell my healthcare provider before taking DAYBUE or DAYBUE STIX?

Before taking DAYBUE or DAYBUE STIX, tell your healthcare provider about all of your medical conditions, including if you:

- have kidney problems.
- are pregnant or plan to become pregnant. It is not known if DAYBUE or DAYBUE STIX will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if DAYBUE or DAYBUE STIX passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby while taking DAYBUE or DAYBUE STIX.

See additional Important Safety Information on page 6.



Choosing between DAYBUE and DAYBUE STIX

- What are the main differences between the two?
- Do they both offer the same benefits?
- What are the different ways they are prepared and stored?



Getting started with DAYBUE

- Do I need to do anything in advance to prepare my loved one to take DAYBUE for the first time?
- Should I let others on my loved one's care team know they are beginning this treatment?
- Are there any support programs or resources available to help us get started?

Important Safety Information (continued)

What should I tell my healthcare provider before taking DAYBUE or DAYBUE STIX? (continued)

Tell your healthcare provider about all of the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking DAYBUE or DAYBUE STIX with certain medicines may affect how the other medicines work and can cause serious side effects.

See additional Important Safety Information on page [6](#).



Taking DAYBUE or DAYBUE STIX

- How is DAYBUE given?
- Can I still give DAYBUE if my loved one has a G-tube or GJ-tube?
- How much DAYBUE will my loved one need to take?
- Are there any administration tips you can give?
- What is the flavor?
- What if we have to travel—how do we take DAYBUE with us?
- What happens if my loved one misses a dose?
- How/where do I store DAYBUE?



Managing possible side effects

- What are the possible side effects of DAYBUE or DAYBUE STIX?
- How many people experienced the side effects?
- If side effects occur, how do I manage them?
- Is there any information about using DAYBUE or DAYBUE STIX in the long term?
- What are signs that I should call my doctor for more help?

Important Safety Information (continued)

What are the possible side effects of DAYBUE or DAYBUE STIX?

- **Diarrhea:** Diarrhea is a common side effect of DAYBUE or DAYBUE STIX that can sometimes be severe. Diarrhea may cause you to lose too much water from your body (dehydration). Before starting treatment with DAYBUE or DAYBUE STIX, stop taking laxatives. Tell your healthcare provider if you have diarrhea while taking DAYBUE or DAYBUE STIX. Your healthcare provider may ask you to increase the amount you drink or take antidiarrheal medicine as needed.

See additional Important Safety Information on page 6.

Indication and Important Safety Information

What is DAYBUE or DAYBUE STIX?

DAYBUE or DAYBUE STIX are prescription medicines used to treat Rett syndrome in adults and children 2 years of age and older. It is not known if DAYBUE or DAYBUE STIX is safe and effective in children under 2 years of age.

What are the possible side effects of DAYBUE or DAYBUE STIX?

- **Diarrhea:** Diarrhea is a common side effect of DAYBUE or DAYBUE STIX that can sometimes be severe. Diarrhea may cause you to lose too much water from your body (dehydration). Before starting treatment with DAYBUE or DAYBUE STIX, stop taking laxatives. Tell your healthcare provider if you have diarrhea while taking DAYBUE or DAYBUE STIX. Your healthcare provider may ask you to increase the amount you drink or take antidiarrheal medicine as needed.
- **Vomiting:** Vomiting is a common side effect of DAYBUE or DAYBUE STIX. Sometimes vomit can get into your lungs (aspiration), which could cause an infection (aspiration pneumonia). Tell your healthcare provider if you have severe vomiting or if vomiting happens often.
- **Weight loss:** DAYBUE or DAYBUE STIX can cause weight loss. Tell your healthcare provider if you notice you are losing weight at any time during treatment with DAYBUE or DAYBUE STIX.
- **The most common side effects of DAYBUE or DAYBUE STIX include diarrhea and vomiting.** Other side effects include fever, seizure, anxiety, decreased appetite, tiredness, and the common cold. These are not all the possible side effects of DAYBUE or DAYBUE STIX. Tell your healthcare provider if you have any side effects that bother you or do not go away.

What should I tell my healthcare provider before taking DAYBUE or DAYBUE STIX?

Before taking DAYBUE or DAYBUE STIX, tell your healthcare provider about all of your medical conditions, including if you:

- have kidney problems.
- are pregnant or plan to become pregnant. It is not known if DAYBUE or DAYBUE STIX will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if DAYBUE or DAYBUE STIX passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby while taking DAYBUE or DAYBUE STIX.

Tell your healthcare provider about all of the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking DAYBUE or DAYBUE STIX with certain medicines may affect how the other medicines work and can cause serious side effects.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call **1-800-FDA-1088**. You can also call Acadia Pharmaceuticals Inc. at **1-844-4ACADIA (1-844-422-2342)**.

DAYBUE is available as an oral solution (200 mg/mL).

DAYBUE STIX for oral solution powder is available in 5,000 mg, 6,000 mg, and 8,000 mg packets.

The risk information provided here is not comprehensive. To learn more, talk with your healthcare provider and read the full [Prescribing Information](#), including [Patient Information](#), also available at DAYBUE.com.